

Tactical Combat Casualty Care Practice Test Questions and Answers

1. What is the first priority in the Care Under Fire phase of TCCC?

- A) Apply pressure dressing to all wounds
- B) Return fire and move the casualty to safety
- C) Start IV fluids immediately
- D) Perform detailed medical assessment

2. Which tourniquet application site is preferred for severe extremity bleeding?

- A) Directly over the wound
- B) 2-3 inches above the wound, high and tight
- C) Below the wound site
- D) Over clothing or gear

3. What is the recommended treatment for tension pneumothorax in tactical settings?

- A) Chest compressions
- B) Needle decompression at the 2nd intercostal space, midclavicular line
- C) High-flow oxygen only
- D) Immediate surgical intervention

4. Which medication is the preferred analgesic for severe pain in TCCC?

- A) Aspirin
- B) Morphine
- C) Ketamine
- D) Ibuprofen

Answers: 1-B 2-B 3-B 4-C

For More Tactical Combat Casualty Care Questions and Answers FREE, Tactical Combat Casualty Care Online Prep Training, Tactical Combat Casualty Care Exam, Tactical Combat Casualty Care Study Guide, Tactical Combat Casualty Care Flashcards, Tactical Combat Casualty Care Quizzes visit:

Tactical Combat Casualty Care Practice Test

Practice Test Geeks © All Rights Reserved